

Log-linear models

The ASTA team

Contents

1	Log-linear models	1
1.1	Model specification	1
1.2	Example	2
1.3	Model specification	2
1.4	Example	2
1.5	Model form	3
1.6	Model comparison	3
1.7	Deviance	4
2	Higher order interaction	4
2.1	Higher order interaction	4
2.2	Bigger example	4
2.3	Test of simple model against the saturated model	6
2.4	Model reduction	7
2.5	Final model and parameter estimates	7
2.6	Predicted values	8
3	Graphical representation	10
3.1	Graphical representation	10
3.2	Interpretation of graphical model	10
3.3	Interpretation of graphical model	12
3.4	Final model of the example	13

1 Log-linear models

1.1 Model specification

- We shall consider the data set `living`, which is a Danish survey on standard of living. The variables are
 - `B` (Housing, Bolig): bad/acceptable/good
 - `H` (Health, Helbred): bad/good
 - `I` (Isolated, Isoleret): yes/no
 - `A` (Anxiety, Angst): yes/no
- `N`: The number of respondents for each combination of the 4 factors above

- We want to order the factors such that the reference is “negative”.

```
living <- read.delim("https://asta.math.aau.dk/datasets?file=living.txt", stringsAsFactors = TRUE)

living$B <- relevel(living$B, "Bad")
living$I <- relevel(living$I, "Yes")
living$A <- relevel(living$A, "Yes")
```

1.2 Example

- At first we study interaction between housing and health. So we aggregate data and only look at the association between B and H without controlling for I and A:

```
BH <- aggregate(N ~ B + H, FUN = sum, data = living)
BH
```

```
##          B     H     N
## 1      Bad  Bad  211
## 2 Acceptable  Bad  327
## 3      Good  Bad 1734
## 4      Bad  Good  145
## 5 Acceptable  Good  211
## 6      Good  Good 1855
```

1.3 Model specification

- Like last time we can write down a model for (the logarithm of) the expected frequencies f_e by using dummy variables.
- We let z_{b1}, z_{b2} and z_{h1} denote the different levels of B and H (the reference level has all dummy variables equal to 0):

$$\log(f_e) = \alpha + \beta_{b1}z_{b1} + \beta_{b2}z_{b2} + \beta_{h1}z_{h1} + \beta_{b1h1}z_{b1}z_{h1} + \beta_{b2h1}z_{b2}z_{h1}.$$

- Note that this time we have included an interaction term, which in this case implies, that we do not assume independence between B and H in the model.
- This model contains all possible terms and there are as many parameters(6) as there are cells(6) in the table. This is called the **saturated** model.

1.4 Example

- We fit the model using `glm`:

```
model <- glm(N ~ B * H, family = poisson, data = BH)
```

- The parameter estimates (of the contrasts, i.e. differences to the reference level (B: Bad, H: Bad) are

```
summary(model)
```

```

## 
## Call:
## glm(formula = N ~ B * H, family = poisson, data = BH)
## 
## Deviance Residuals:
## [1] 0 0 0 0 0 0
## 
## Coefficients:
##              Estimate Std. Error z value Pr(>|z|)
## (Intercept) 5.35186   0.06884 77.740 < 2e-16 ***
## BAcceptable 0.43810   0.08830  4.961 7.00e-07 ***
## BGood        2.10633   0.07291 28.889 < 2e-16 ***
## HGood        -0.37512   0.10787 -3.478 0.000506 ***
## BAcceptable:HGood -0.06298   0.13940 -0.452 0.651438
## BGood:HGood    0.44258   0.11292  3.919 8.88e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## 
## (Dispersion parameter for poisson family taken to be 1)
## 
## Null deviance: 4.2103e+03 on 5 degrees of freedom
## Residual deviance: -2.9532e-14 on 0 degrees of freedom
## AIC: 59.485
## 
## Number of Fisher Scoring iterations: 2

```

- The combination Good:Good increases the response, i.e. an over representation of people with good housing conditions and good health.

1.5 Model form

- Log-linear models quickly become cumbersome to write down.
- For log-linear models the structure of the model is often more interesting than the value of the parameters.
- Therefore we typically just use the model form

$$B + H + B : H$$

where $B + H$ are the main effects and $B : H$ means that we have interaction between B and H .

- We shall stick to the **hierarchical principle**, which means that if $B : H$ is included then we must include $B + H$. For that reason we use the shorthand notation

$$B * H = B + H + B : H$$

1.6 Model comparison

- If we suggest a simpler model than the saturated model it will always provide a poorer fit to the given data.
- We need to find a model which is as simple (few parameters) as possible but which at the same time fits the data well.
- Last time we used the χ^2 statistic to judge whether the model $B + H$ (independence between B and H) was good enough compared to the saturated model $B * H$. This is only possible for models with two variables.

- More generally we shall consider the **Deviance of a model**, which is - approximately - given by

$$\sum \frac{(\text{observed} - \text{expected})^2}{\text{expected}}$$

1.7 Deviance

- We look at the output from the function `drop1`:

```
drop1(model, test = "Chisq")

## Single term deletions
##
## Model:
## N ~ B * H
##      Df Deviance    AIC    LRT Pr(>Chi)
## <none>     0.000 59.485
## B:H      2   40.766 96.251 40.766 1.405e-09 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

- The deviance for the saturated model (<none>) is zero as observed=expected.
- Deviance is increased by 40.8, when we remove the interaction B:H.
- Removing B:H corresponds to the null hypothesis $H_0 : \beta_{b1h1} = \beta_{b2h1} = 0$.
- The deviance is compared with a $\chi^2(Df = 2)$ distribution to determine whether it is significant.
- Since the p-value is 1.4×10^{-9} the interaction is significant and cannot be left out.

2 Higher order interaction

2.1 Higher order interaction

- We shall consider models with interaction between more than two variables.
- E.g. there may be a combined effect of adding water, fertilizer and light to a plant at the same time, i.e. the effect of adding water and fertilizer depends on whether the light is on.
- We call this a **3-way interaction**.
- We shall still respect the hierarchical principle:
 - If we include a 3-way interaction, then we must include all main effects and 2-way interactions of the 3 variables.
- Again we use short hand notation

$$B * H * A = B + H + A + B : H + B : A + H : A + B : H : A$$

- Similar considerations hold for 4-way interactions like $B * H * A * I$, etc.

2.2 Bigger example

- We fit the saturated model to the full dataset and look at the estimated parameters (only the last half of the values are printed to save space):

```

satmodel <- glm(N ~ B * H * A * I, family = poisson, data = living)
summary(satmodel)

##
## Call:
## glm(formula = N ~ B * H * A * I, family = poisson, data = living)
##
## Deviance Residuals:
## [1] 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
## [24] 0
##
## Coefficients:
##                               Estimate Std. Error z value Pr(>|z|)
## (Intercept)           2.079e+00 3.536e-01 5.882 4.06e-09 ***
## BAcceptable          5.596e-01 4.432e-01 1.263 0.206710
## BGood                1.417e+00 3.941e-01 3.596 0.000323 ***
## HGood               -2.438e+01 4.225e+04 -0.001 0.999540
## ANo                  4.855e-01 4.494e-01 1.080 0.279943
## INo                 1.749e+00 3.831e-01 4.566 4.96e-06 ***
## BAcceptable:HGood   2.284e+01 4.225e+04 0.001 0.999569
## BGood:HGood          2.268e+01 4.225e+04 0.001 0.999572
## BAcceptable:ANo     5.349e-02 5.613e-01 0.095 0.924076
## BGood:ANo            -3.077e-02 5.015e-01 -0.061 0.951069
## HGood:ANo            2.343e+01 4.225e+04 0.001 0.999558
## BAcceptable:INo     6.192e-03 4.801e-01 0.013 0.989710
## BGood:INo            5.473e-01 4.244e-01 1.290 0.197159
## HGood:INo            2.405e+01 4.225e+04 0.001 0.999546
## ANo:INo              6.557e-01 4.802e-01 1.365 0.172142
## BAcceptable:HGood:ANo -2.345e+01 4.225e+04 -0.001 0.999557
## BGood:HGood:ANo      -2.254e+01 4.225e+04 -0.001 0.999574
## BAcceptable:HGood:INo -2.303e+01 4.225e+04 -0.001 0.999565
## BGood:HGood:INo      -2.267e+01 4.225e+04 -0.001 0.999572
## BAcceptable:ANo:INo  -2.516e-01 6.007e-01 -0.419 0.675370
## BGood:ANo:INo         2.827e-01 5.329e-01 0.531 0.595707
## HGood:ANo:INo         -2.339e+01 4.225e+04 -0.001 0.999558
## BAcceptable:HGood:ANo:INo 2.365e+01 4.225e+04 0.001 0.999553
## BGood:HGood:ANo:INo   2.301e+01 4.225e+04 0.001 0.999565
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
## Null deviance: 1.0965e+04 on 23 degrees of freedom
## Residual deviance: 4.1199e-10 on 0 degrees of freedom
## AIC: 179.1
##
## Number of Fisher Scoring iterations: 20

```

- We see that the four-way interaction and all three-way interactions look like they could be left out.
- However, the p-values are related to removing one and leaving everything else in the model so we have to remove terms of the model one at a time and check against the new model.
- This can be done by successive use of `drop1` and `update` as explained in the following.

```

drop1(satmodel, test = "Chi")

## Single term deletions
##
## Model:
## N ~ B * H * A * I
##          Df Deviance    AIC    LRT Pr(>Chi)
## <none>      0.0000 179.10
## B:H:A:I  2    3.5112 178.61 3.5112   0.1728

```

- The output of `drop1` reveals that the four-way interaction is insignificant and we remove it and save the updated model like this:

```

reducedmodel <- update(satmodel, .~.-B:H:A:I)

```

- We use `drop1` again:

```

drop1(reducedmodel, test = "Chi")

## Single term deletions
##
## Model:
## N ~ B + H + A + I + B:H + B:A + H:A + B:I + H:I + A:I + B:H:A +
##           B:H:I + B:A:I + H:A:I
##          Df Deviance    AIC    LRT Pr(>Chi)
## <none>      3.5112 178.61
## B:H:A  2    7.1419 178.24 3.6307   0.1628
## B:H:I  2    3.5316 174.63 0.0205   0.9898
## B:A:I  2    5.5332 176.63 2.0220   0.3639
## H:A:I  1    4.5857 177.68 1.0745   0.2999

```

- We see `B:H:I` could be removed and use `update` again:

```

reducedmodel <- update(reducedmodel, .~.-B:H:I)

```

- We can continue this way as long as we like.
- If instead we have a specific simple model in mind, we can define this model and test it against the saturated model with `anova` as shown in the following.

2.3 Test of simple model against the saturated model

- We fit the simpler model to the full dataset:

```

simplemodel <- glm(N ~ B*H + B*A + B*I + H*A + H*I + I*A, family = poisson, data = living)

```

- We compare the two models using `anova`:

```

anova(simplemodel, satmodel, test = "Chisq")

## Analysis of Deviance Table
##
## Model 1: N ~ B * H + B * A + B * I + H * A + H * I + I * A
## Model 2: N ~ B * H * A * I
##   Resid. Df Resid. Dev Df Deviance Pr(>Chi)
## 1          9    10.697
## 2          0    0.000  9    10.697  0.2971

```

- The deviance between the two models is 10.697 with $df = 9$, which has a p-value of 29.7%. So we prefer the simpler model without four- and 3-way interactions.

2.4 Model reduction

- Let us check whether we can make further model reductions, where we again use the function `drop1`.

```

drop1(simplemodel, test = "Chisq")

## Single term deletions
##
## Model:
## N ~ B * H + B * A + B * I + H * A + H * I + I * A
##           Df Deviance    AIC    LRT  Pr(>Chi)
## <none>     10.697 171.79
## B:H      2    38.170 195.27 27.474 1.082e-06 ***
## B:A      2    37.912 195.01 27.215 1.231e-06 ***
## B:I      2    35.245 192.34 24.548 4.671e-06 ***
## H:A      1    41.947 201.04 31.250 2.268e-08 ***
## H:I      1    56.048 215.14 45.352 1.646e-11 ***
## A:I      1    26.449 185.54 15.753 7.218e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

- Conclusion: All of the pairwise interaction terms are significant.

2.5 Final model and parameter estimates

- In conclusion our final model is `simplemodel`.

```

summary(simplemodel)

##
## Call:
## glm(formula = N ~ B * H + B * A + B * I + H * A + H * I + I *
##       A, family = poisson, data = living)
##
## Deviance Residuals:
##       Min        1Q    Median        3Q        Max

```

```

## -1.72955 -0.48421 -0.00248  0.31663  1.23187
##
## Coefficients:
##                               Estimate Std. Error z value Pr(>|z|)
## (Intercept)            2.17042   0.22962  9.452 < 2e-16 ***
## BAcceptable           0.65323   0.26323  2.482 0.013081 *
## BGood                 1.13785   0.23365  4.870 1.12e-06 ***
## HGood                -1.76785   0.21028 -8.407 < 2e-16 ***
## ANo                  0.35067   0.19249  1.822 0.068488 .
## INo                  1.74803   0.23116  7.562 3.97e-14 ***
## BAcceptable:HGood   -0.04121   0.14119 -0.292 0.770370
## BGood:HGood          0.37773   0.11441  3.302 0.000961 ***
## BAcceptable:ANo     -0.12759   0.15834 -0.806 0.420355
## BGood:ANo             0.39413   0.13265  2.971 0.002966 **
## BAcceptable:INo     -0.14021   0.25873 -0.542 0.587878
## BGood:INo             0.72038   0.22799  3.160 0.001579 **
## HGood:ANo             0.44076   0.07938  5.552 2.82e-08 ***
## HGood:INo             1.12352   0.17982  6.248 4.16e-10 ***
## ANo:INo              0.66875   0.16277  4.109 3.98e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
## Null deviance: 10964.662 on 23 degrees of freedom
## Residual deviance: 10.697 on 9 degrees of freedom
## AIC: 171.79
##
## Number of Fisher Scoring iterations: 4

```

- We see that all significant interactions are positive - as expected(why?).
- (Intercept) is log of the expected number, when all factors are “bad”. So the expected number is $\exp(2.17) = 8.76$, whereas the observed number is 8.

2.6 Predicted values

- What is the expected number of people without anxiety (A), with acceptable housing (B), good health (H) that are isolated (I)?

```

summary(simplemodel)

##
## Call:
## glm(formula = N ~ B * H + B * A + B * I + H * A + H * I + I *
##      A, family = poisson, data = living)
##
## Deviance Residuals:
##      Min        1Q        Median         3Q        Max
## -1.72955 -0.48421 -0.00248  0.31663  1.23187
##
## Coefficients:
##                               Estimate Std. Error z value Pr(>|z|)

```

```

## (Intercept) 2.17042 0.22962 9.452 < 2e-16 ***
## BAcceptable 0.65323 0.26323 2.482 0.013081 *
## BGood 1.13785 0.23365 4.870 1.12e-06 ***
## HGood -1.76785 0.21028 -8.407 < 2e-16 ***
## ANo 0.35067 0.19249 1.822 0.068488 .
## INo 1.74803 0.23116 7.562 3.97e-14 ***
## BAcceptable:HGood -0.04121 0.14119 -0.292 0.770370
## BGood:HGood 0.37773 0.11441 3.302 0.000961 ***
## BAcceptable:ANo -0.12759 0.15834 -0.806 0.420355
## BGood:ANo 0.39413 0.13265 2.971 0.002966 **
## BAcceptable:INo -0.14021 0.25873 -0.542 0.587878
## BGood:INo 0.72038 0.22799 3.160 0.001579 **
## HGood:ANo 0.44076 0.07938 5.552 2.82e-08 ***
## HGood:INo 1.12352 0.17982 6.248 4.16e-10 ***
## ANo:INo 0.66875 0.16277 4.109 3.98e-05 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for poisson family taken to be 1)
##
## Null deviance: 10964.662 on 23 degrees of freedom
## Residual deviance: 10.697 on 9 degrees of freedom
## AIC: 171.79
##
## Number of Fisher Scoring iterations: 4

```

- Logarithm of the expected:

$$2.17042 + 0.65323 - 1.76785 + 0.35067 - 0.04121 - 0.12759 + 0.44076 = 1.67843$$

- so the expected number is $\exp(1.67843) = 5.36$, whereas the observed number is 5.

- We can of course make **R** calculate all the table values expected by the model and add them to the data:

```

living$expected <- fitted(simplemodel)
living[1:12,]

```

```


##      B   H   I   A   N   expected
## 1 Bad Good Yes No  5  3.300261
## 2 Bad Good No  No 107 113.784094
## 3 Bad Good Yes Yes  0  1.495663
## 4 Bad Good No Yes 33  26.419981
## 5 Bad Bad Yes No 13  12.442145
## 6 Bad Bad No  No 144 139.473499
## 7 Bad Bad Yes Yes  8  8.761930
## 8 Bad Bad No Yes 46  50.322426
## 9 Acceptable Good Yes No  5  5.357140
## 10 Acceptable Good No  No 155 160.536219
## 11 Acceptable Good Yes Yes  3  2.758237
## 12 Acceptable Good No Yes 48  42.348403

```

3 Graphical representation


3.1 Graphical representation

- We make a graphical representation by
 - drawing a circle for each variable.
 - connecting variables which enter the same model term.
- Example: Assume the model is
$$A * B + B * H * I$$
- Then the graphical representation is




3.2 Interpretation of graphical model

- **Independence:** If A enters in the model formula, but A doesn't enter in any other terms (e.g. $A * B$, $A * H$, etc.), then A is independent of the other variables.
- E.g. $A + B * H + B * I$




- **Explained association.** If B and H are "connected" via other terms, but don't enter the same term, then the association is explained by other variables. I.e. the model cannot include e.g. $B * H$, $B * H * A$ or $A * B * H * I$.
- E.g. $B * I + A * H * I$




3.3 Interpretation of graphical model

- **Homogeneous association:** If $A * H$ enters the model, but $A * H$ doesn't enter more complicated terms, then the association between A and H is homogeneous.
 - I.e. the model cannot contain $A * H * I$, $A * B * H$ or $A * B * H * I$.
 - E.g. $A * H + A * I * B + B * H$




- **Heterogeneous association:** If $A * H$ enter the model as a part of a more complicated term, then the association between A and H is heterogeneous.
 - I.e. the model *must* contain $A * H * I$, $A * B * H$ or $A * B * H * I$.
 - E.g. $A * B * H + A * I * B$



3.4 Final model of the example

- In the example the final model was:

$$B * I + H * I + I * A + B * H + B * A + H * A$$



- We can directly see from the graph, that:
 - we don't have any independent variables since all variables are connected
 - we do not have an explained association.
- From the formula we have homogeneous association between all pairs of variables.