15.WAIS

```
library(mosaic)
```

Exercise 15.WAIS.

Referring to exercise 15.3 in Agresti, one of the WAIS subtests called *picture completion*, asks questions about 20 pictures that have one vital detail missing. It is considered a test of attention to fine detail.

The observations for 20 subjects on (x,y), where x=picture completion score(0-20) and y=symptoms of senility(1=yes):

```
pictSen <-
read.table("https://asta.math.aau.dk/datasets?file=pictSen.csv",header=T)
pictSen</pre>
```

##		pictComplScore	senility
##	1	7	1
##	2	5	1
##	3	3	1
##	4	8	1
##	5	1	1
##	6	2	1
##	7	9	1
##	8	3	1
##	9	6	1
##	10	4	1
##	11	6	0
##	12	9	0
##	13	7	0
##	14	7	0
##	15	10	0
##	16	12	0
##	17	14	0
##	18	8	0
##	19	8	0
##	20	11	0

• Assume a simple logistic model and estimate the logistic regression equation.

```
mod <- glm(senility ~ pictComplScore, data = pictSen, family = binomial)
coef(summary(mod))</pre>
```

```
## Estimate Std. Error z value Pr(>|z|)
## (Intercept) 5.426569 2.6279764 2.064923 0.03893027
## pictComplScore -0.772115 0.3578408 -2.157706 0.03095074
```

- Why is there a significant effect of picture completion on symptoms?
 - p-value is small.

- Estimate the probability that symptoms are present when (i) x=0, (ii) x=20.
 - (i) ilogit(5.4266)=0.9956212.
 - (ii) ilogit(5.4266-0.7721*20)= 4.4704128×10^{-5} .
- Over what range of x-scores is the estimated probability of senility greater than 50%?
 - It is greater than 50% for any score below:

```
a <- 5.4266
b <- -0.7721
-a/b
```

[1] 7.028364

- Estimate the effect of a one-unit increase in picture completion on the odds of senility symptoms.
 - For every one-unit increase in picture completion the odds decrease approximately 50%:

```
exp(b)-1
```

[1] -0.5379582

- Provide a 95% confidence interval of this effect.
 - From the model output we know the std. error of the effect is 0.358. An approximate 95% confidence interval is:

```
\exp(b+c(-2, 2)*0.358)-1
```

[1] -0.77419873 -0.05455541