

Module 7: Solutions for supplementary exercises

Exercise: Potential rejection sampling problems

First try to answer the following questions without using the computer – then reuse the code from the supplementary slides to check your answer:

- Suppose we could not easily determine M and hence had to make a conservative choice; say $M = 100$ or $M = 500$ in this context.
 1. Which effect will that have on the number of accepted samples? **The acceptance rate goes down.**

```
f0 <- function(x, a=.4, b=.08){exp(a * (x - a)^2 - b * x^4)}
N <- 10000
M <- 500
y <- runif(N, -4, 4)
p_accept <- f0(y)/(M*dunif(y, -4, 4))
u <- runif(N, 0, 1)
keep <- u<p_accept
mean(keep)
```


```
## [1] 0.0153
```

2. How would you have to compensate for a too large value of M if you want a given number of samples from the target distribution?
****Increase the number of proposals, which will increase the overall computation time.****

- What happens if you do not choose M large enough (e.g. $M = 10$ in our example)? **Then you sample from the wrong distribution.**

```
M <- 10
y <- runif(N, -4, 4)
p_accept <- f0(y)/(M*dunif(y, -4, 4))
u <- runif(N, 0, 1)
keep <- u<p_accept
hist(y[keep], prob = TRUE, col = "gray", ylim = c(0, .38))
norm_const <- integrate(f0, -4, 4)$value
curve(f0(x)/norm_const, col = "red", add = TRUE)
```

Histogram of y[keep]



- What would be the effect of using a uniform proposal distribution on $[-10, 10]$? **Acceptance rate goes down since proposals outside $[-4, 4]$ always are rejected.**

```
M <- 3.1/20
y <- runif(N, -20, 20)
p_accept <- f0(y)/(M*dunif(y, -20, 20))
u <- runif(N, 0, 1)
keep <- u < p_accept
mean(keep)
```

```
## [1] 0.17
```

- What happens if the proposal distribution is a standard normal distribution (i.e. mean zero and standard deviation 1)? Hints:
 1. You can use `dnorm()` for the normal density.
 2. You may have to create a sequence `x <- seq(-4, 4, by = 0.01)` to numerically evaluate the bound M relating $f_0(x)$ and $dnorm(x)$.

The upper bound is higher and we get lower acceptance rates. I.e. we have to sample for longer time to obtain the number of target samples we want.

```
x <- seq(-4, 4, by = 0.01)
M <- max(f0(x)/dnorm(x))
y <- rnorm(N)
p_accept <- f0(y)/(M*dnorm(y))
u <- runif(N, 0, 1)
keep <- u < p_accept
mean(keep)
```

```
## [1] 0.1063
```

Exercise: Improving the proposal distribution

If $f(x)$, $x \in [0, 1]$ is a pdf on $[0, 1]$ then for $a > 0$, $1/a \cdot f(x/a)$, $x \in [0, a]$ is a pdf on $[0, a]$. Furthermore, a pdf on $[b, a + b]$ can be obtained by simple translation.

- Based on these facts how can a beta distribution $\text{Be}(\alpha, \beta)$ indirectly be used as the proposal distribution for our example? -Implement the rejection sampling algorithm using $\text{Be}(2.5, 3.5)$ transformed to $[-4.1, 4.1]$ (but with M determined on $[-4, 4]$).
- Check with a histogram that you are sampling the correct distribution.
- Find the acceptance rate.

For $X \sim \text{Be}(2.5, 3.5)$ transform it to the interval $I = [-4.1, 4.1]$ by $Y = 8.2 \cdot X - 4.1$. Then the density for Y is $f((y + 4.1)/8.2)/8.2$ for x in I and zero otherwise, where f is the original Beta density on $[0, 1]$:

```
x <- seq(-4, 4, by = 0.01)
g <- function(y, a = 2.5, b = 3.5){x <- (y+4.1)/8.2; dbeta(x, a, b)/8.2}
M <- max(f0(x)/g(x))
y <- 8.2*rbeta(N, 2.5, 3.5) - 4.1
p_accept <- f0(y)/(M*g(y, 2.5, 3.5))
u <- runif(N, 0, 1)
keep <- u < p_accept
mean(keep)
```

```
## [1] 0.5674
```

```
hist(y[keep], prob = TRUE, col = "gray", ylim = c(0, .38))
curve(f0(x)/norm_const, col = "red", add = TRUE)
```

